

WITHDRAWAL

Benzodiazepines  
Gabapentin  
Ketamine  
Clonidine  
Tizanidine  
Penobarbital  
Dexmedetomidine

Recommended Medication for Tranq withdrawal:

Individuals can differentiate symptoms of xylazine withdrawal from opiate withdrawal when being properly treated with OAT like: Suboxone, Kadian, Hydromorphone

WITHDRAWAL

Anxiety  
Irritability  
Insomnia  
Agitation  
Risk of seizure

Xylazine withdrawal is similar to benzo withdrawal.

Opioid withdrawal should be treated early, with liberal use of medications for opioid use disorder or opioid pain management in order to mitigate any pain and discomfort that could further exacerbate the manifestations of xylazine withdrawal or lead to discharges against medical advice.

WITHDRAWAL

Anxiety  
Irritability  
Insomnia  
Agitation  
Risk of seizure

Xylazine withdrawal is similar to benzo withdrawal.

Opioid withdrawal should be treated early, with liberal use of medications for opioid use disorder or opioid pain management in order to mitigate any pain and discomfort that could further exacerbate the manifestations of xylazine withdrawal or lead to discharges against medical advice.

WITHDRAWAL

Anxiety  
Irritability  
Insomnia  
Agitation  
Risk of seizure

Xylazine withdrawal is similar to benzo withdrawal.

Opioid withdrawal should be treated early, with liberal use of medications for opioid use disorder or opioid pain management in order to mitigate any pain and discomfort that could further exacerbate the manifestations of xylazine withdrawal or lead to discharges against medical advice.